

ANATOMIC VARIATIONS OF PARANASAL SINUSES IN PATIENTS WITH CHRONIC SINUSITIS AND THEIR CORRELATION WITH CT SCAN STAGING

M. H. Baradaranfar^{1,2}, M. Labibi³

1) Department of Otolaryngology and Head and Neck Surgery, School of Medicine, Yazd University of Medical Sciences, Yazd, Iran

2) Otorhinolaryngology Research Center, Medical Sciences/University of Tehran, Tehran, Iran

3) Department of Otolaryngology and Head and Neck Surgery, School of Medicine, Chaharmahal-Bakhtiari University of Medical Sciences, Chaharmahal-Bakhtiari, Iran

Abstract- The presence of anatomic variations of paranasal sinuses must be noted in order to attain a full understanding and accurate diagnosis of chronic sinusitis. The frequency of anatomic variations in chronic sinusitis is different in various studies. The purpose of this study is to assess the frequency of anatomic variations in chronic sinusitis patients admitted to Otolaryngology ward of Shahid Sadoughi hospital in the year 2004 and underwent functional endoscopic sinus surgery. Based on preoperative paranasal sinus CT scan the extent of patients diseases on a scale of 0-IV and the presence of anatomic variations was determined. A total of 120 patients were included. We found Agger Nasi cells in 43(36%), Haller cells in 5 (4.17%) conchae bullosa in 15(12.5%), paradoxical middle turbinate in 2 (1.67%) and septal deviation in 54 patients (45%). Patients with Agger Nasi (P.value : 0.015) and conchae bullosa (P.Value : 0.024) demonstrated significantly higher scores on sinus CT scan compared with patients without Agger Nasi cells and conchae bullosa respectively. Three months postoperatively we observed significantly less resolution of nasal congestion (P.Value : 0.006) in patients with septal deviation than patients without it. No significant difference was seen among patients with and without other anatomic variations with regard to resolution of rhinorrhea and nasal congestion three months postoperatively.

© 2007 Tehran University of Medical Sciences. All rights reserved.

Acta Medica Iranica, 45(6): 477-480; 2007

Key Words : Paranasal sinuse , chronic sinusitis, CT Scan staging

INTRODUCTION

Although the anatomy of nasal cavity and paranasal sinuses differ significantly from patient to patient, certain distinct variations are found most frequently among the general population. The presence of anatomic variations must be noted in

order to attain a full understanding of the individual patient as well as to develop an accurate diagnosis (1,2). The purpose of this study was to assess the frequency of anatomic variation of nasal cavity and paranasal sinuses in patients older than 16 years old with chronic sinusitis refractory to medical therapy who are candidates for functional endoscopic sinus surgery. Reported frequency of anatomic variations in patient with chronic sinusitis is as follow: Agger Nasi cellss in 15% , Haller cells in 7% , conchae bullosa in 30% , paradoxical middle turbinate in 24% and septal deviation in 21% of patients (3).Endoscopic surgery of the chronic sinusitis needs an accurate evaluation of disease and paranasal

Received: 27 Feb. 2006, Revised: 22 Apr. 2006, Accepted: 21 Aug. 2006

* Corresponding Author:

Mohammad Hossein Baradaranfar, Department of Otolaryngology and Head and Neck Surgery, School of Medicine, Yazd University of Medical Sciences, Yazd, Iran Fax: + Tel: +98 351 8224000
98 351 8224100

E-mail: baradaranf@ yahoo.com

anatomic variations. After identification of these variations, functional endoscopic sinus surgery with usually minimal invasive operations can provide dramatic relief of chronic sinusitis symptoms (4)

Some claim that the mere presence of anatomic variations causes narrowing or obstruction of the osteomeatal channels, thereby hampering the normal airflow and mucociliary clearance of the sinuses while others contend that whether an anatomic variation is a factor in development of chronic sinusitis depends on its size, location or the amount of mucosal contact caused by the variation (2,5).

MATERIALS AND METHODS

From January to November 2004, patients (older than 16 years) with chronic sinusitis with symptoms such as purulent rhinorrhea, nasal congestion, headache, facial pain or pressure and cough persisting beyond rhinorrhea and congestion persisting beyond 12 weeks who had not demonstrated symptom resolution despite 3-6 weeks antibiotic therapy, intranasal or systemic steroids, decongestants and nasal saline irrigation were included in the study. All patients without subjective relief of symptoms underwent examination of nose and middle meatus with 30- degree telescope to obtain clinical information regarding presence of purulent secretions and polypoid mucosa in the middle meatus and those with positive findings underwent coronal CT scan of pronasal sinuses in preparation for operation. A total of 120 patients (79 male and 41 female) aged 17 to 79 years (mean age of 34.46) with chronic sinusitis were admitted to Otolaryngology ward of Shahid Sadoughi hospital and underwent functional endoscopic sinus surgery in the operating room. The extent of patient disease on a scale of 0-IV (sinus CT staging system) (6) and

presence of anatomic variations such as Agger Nasi cells, Haller cells , conchae bullosa, paradoxical middle turbinate and nasal septal deviation were determined based on paranasal sinus coronal CT scan and intraoperative findings.

Stage I: All unilateral disease or anatomic abnormality

Stage II: Bilateral disease limited to ethmoid or maxillary sinus.

Stage III: Bilateral disease with involvement of at least one sphenoid or frontal sinus.

Stage IV: Pansinusitis

Three months postoperatively all patients were evaluated for resolution of rhinorrhea and nasal congestion. Resolution of rhinorrhea was determined by anterior rhinoscopy and oral examination for presence of posterior nasal drip. The degree of congestion relief(slight, relative and complete) was based on patients own description of nasal patency compared with preoperative state and examination of nasal airflow with a piece of cotton.

RESULTS

A total of 120 patients were enrolled in the study. We found Agger Nasi cells in 43(36%) , Haller cells in 5(4.17%) , conchae bullosa in 15(12.5%) , paradoxical middle turbinate in 2(1.67%) and nasal septal deviation in 54 patients (45%).

According to paranasal sinus CT scan and operative findings we noticed stage I, II, III and IV in 9,24,50 and 37 patients respectively.

Patients with Agger Nasi (p= 0.015) and conchae bullosa (p= 0.024) demonstrated significantly higher scores on sinus CT scan compared with patients without Agger Nasi and conchae bullosa respectively (Table 1 and 2).

Table.1. The frequency of Agger Nasi cells in chronic sinusitis according to sinus CT staging

Agger Nasi		Stage				Total
		I	II	III	IV	
No	Count	9	17	25	26	77
	Percent	11.7%	22.1%	32.5%	33.8%	100%
Yes	Count	0	7	25	11	43
	Percent	0%	16.3	58.1%	25.6%	100%
Total	Count	9	24	50	37	120
	Percent	7.5%	20.0%	41.7%	30.8%	100%

p = 0.015

Table 2. The frequency of conchae bullosa in chronic sinusitis according to sinus CT staging

Conchae Bullosa		Stage				Total
		I	II	III	IV	
No	Count	9	23	43	30	105
	Percent	8.6%	27%	35.8%	28.6%	100%
Yes	Count	0	1	7	7	15
	Percent	0%	6.7%	46.7%	46.7%	100%
Total	Count	9	24	50	37	120
	Percent	7.5%	20.0%	41.7%	30.8%	100%

p = 0.024

No significant difference was seen among patients with and without Haller cells, paradoxical middle turbinate and nasal septal deviation with regard to sinus CT staging system.

Three months postoperatively, we observed significantly less resolution of nasal congestion in patients with nasal septal deviation (p= 0.006) than patients without septal deviation (Table 3).

No significant difference was seen among patients with and without Agger Nasi cells, conchae bullosa and paradoxical middle turbinate with regard to postoperative resolution of nasal congestion.

Postoperative resolution of rhinorrhea was not significantly different among patients with and without the five abovementioned anatomic variations.

DISCUSSION

In this study the frequency of anatomic variations of paranasal sinuses in a group of Iranian patients with chronic sinusitis is determined.

Our findings correspond with results of other studies (2,7,8) with regard to frequency of Agger Nasi cells, Haller cells and septal deviation but the frequency of conchae bullosa and paradoxical middle turbinate in our patients was less.

Some authors define concha bullosa as any degree of middle turbinate pneumatization (2) but in our study it was defined as pneumatization of both vertical lamellar and inferior bulbous portions of middle turbinate with obstruction of middle meatus which explains for its frequency of 12.5% in our patients compared with frequency of 35%(9), 30% (10) and 28% (11) in other studies.

Lloyd reported that obtaining an accurate image of paradoxically curved middle turbinate was related to the level of the coronal CT scan. Some patients in his study showed paradoxical curves anteriorly and normal curves posteriorly (2).

The abovementioned statement explains for low frequency of paradoxical middle turbinate in our patients with inadequate anterior coronal cuts in paranasal sinus CT scan.

Table 3. The frequency of septal deviation in chronic sinusitis according to postoperative resolution of nasal congestion

Septal deviation		Postoperative resolution of nasal congestion			Total
		Slight	Relative	Complete	
No	Count	0	11	55	66
	Percent	0%	16.7%	83.3%	100%
Yes	Count	1	22	31	54
	Percent	1.9%	40.7%	57.4%	100%
Total	Count	1	33	86	120
	Percent	0.8%	27.5%	71.7%	100%

p = 0.006

Anatomic variations of paranasal sinuses

Table 1 and 2 demonstrate significantly higher scores (stage III and IV) in patients with Agger Nasi and Concha bullosa which signifies their role in frontal sinus obstruction and obstruction of middle meatus and infundibulum respectively (12,13).

Three months postoperatively, less resolution of nasal congestion was seen significantly in patients with septal deviation, so we recommend performing septoplasty in combination with functional endoscopic sinus surgery.

REFERENCES

1. Aygun N, Zinreich SJ, Radiology of the Nasal Cavity and Paranasal sinuses. Cummings CW et al, Cummings Otolaryngology Head & Neck Surgery. 4th ed. Elsevier Mosby; 2005; P. 1159-1160.
2. Kennedy DW, Bolger, WE. Zinreich SJ : Disease of the Sinuses, Diagnosis and Management, B.C. Decker Inc. Hamilton. London. 2001; P. 19-24.
3. Hosemann WG, Weber RK, Keerl RE, Lund VJ. Minimally Invasive Endonasal Sinus Surgery, Thieme, Stuttgart, New York. 2000; P. 3.
4. Arslan H, Aydinlioglu A, Egeli E, Anatomic variations of the paranasal sinuses: CT examination for endoscopic sinus surgery. International Journal of ORL and HNS. 2004 Feb; :39-48
5. TINGT-KUANG CHAO. Uncommon anatomic variations in patients with chronic paranasal sinusitis otolaryngology- Head & Neck Surgery. 2005 Feb; 132(2):221-5
6. Montgomery WW. Surgery of the Upper Respiratory System, William & Wilkins: 3rd ed. 1996; P. 193-194.
7. Eryilmaz A, Gocer C, Dursun E. The incidence of anatomic variations and sinus opacities in patients with chronic sinonasal symptoms, Kulak Burun Bogaz Ihtis Derg. 2004; 13(5-6):116-121.
8. Krzeski A, Tomaszewska E, Jakubczyki. Anatomic variations of the lateral nasal wall in the computed tomography scan of patients with chronic rhinosinusitis. Am J Rhinol. 2001 Nov-Dec; 15(6): 371-375.
9. Bolger WE, Butzin CA, Parsons DS. Paranasal sinus bony anatomic variation and mucosal abnormalities: CT analysis for endoscopic sinus surgery. Laryngoscope. 1991; 101:56-64.
10. Wanamaker HH. Role of Haller's cell in headache and sinus disease: a case report. Otolaryngol Head Neck Surg. 1996; 114: 324-327.
11. Tonia A, Baba S. Anatomic variations of the bone in sinonasal CT. Acta Otolaryngol suppl (stockh). 1996; 525:9-13.
12. Palmer JN, Kennedy DW, Revision Endoscopic Sinus Surgery. Cummings Otolaryngology Head & Neck Surgery. 4th ed. Elsevier Mosby: 2005; P. 1268-1269.
13. Sivasli E, Sirikci A, Bayazyt YA. Anatomic variations of paranasal sinuses area in patients with chronic sinusitis. Surg Radiol Anat. 2003 Feb; 24(6):400-405