

Association of Maternal Serum C- Reactive Protein Levels with Severity of Preeclampsia

Fatemeh Mirzaie^{1*}, Fatemeh Rahimi-Shorbat², and Amir Hossian Kazeroni³

¹ Department of Obstetric and Gynecology, School of Medicine, Kerman University of Medical Sciences, Kerman, Iran

² Department of Fetal Maternal Medicine, School of Medicine, Tehran University of Medical Sciences, Tehran, Iran

³ Department of General Practitioner, School of Medicine, Kerman University of Medical Sciences, Kerman, Iran

Received: 3 Sep. 2007; Received in revised form: 6 Apr. 2008; Accepted: 14 May 2008

Abstract- The aim of this study was to investigate C-reactive protein (CRP) level in preeclampsia (PE) and its association with the severity of the disease. This cross-sectional study included 43 women with mild PE, 43 women with severe PE, and 43 healthy pregnant. They were selected in the third trimester of pregnancy in the Afzalipour Hospital, Kerman, Iran, from March 2006 to March 2007. Mean diastolic pressure and level of proteinuria were used as indicators of the severity of the disease. The results were analyzed by t-test and spearman's rank correlation coefficient. Hemoglobin, aspartate and alanine transaminase, creatinine and urine protein excretion, serum CRP, and alkaline phosphatase were higher in women with PE. There were significant correlations between serum CRP levels and diastolic blood pressure ($r = 0.5$, $P = 0$), urinary protein excretion ($r = 0.5$, $P = 0$), creatinine ($r = 0.2$, $P = 0.003$), spartate transaminase ($r = 0.3$, $P = 0$), alanine transaminase ($r = 0.2$, $P = 0.006$), and Hemoglobin ($r = 0.2$, $P = 0.001$). There were a negative correlation between serum CRP and weight of the new born ($r = -0.09$, $P = 0.01$) and gestational age in the time of delivery ($r = -0.07$, $P = 0$). We showed higher levels of CRP in women with PE. Elevated serum levels of CRP in PE women are, thus, correlated with severity of disease.

© 2009 Tehran University of Medical Sciences. All rights reserved.

Acta Medica Iranica 2009; 47(4): 293-296.

Key words: C - reactive protein, severity of disease, preeclampsia

Introduction

Preeclampsia (PE) develops in 4-5% of human pregnancies. It is characterized by an elevated blood pressure and proteinuria and develops after 20 weeks of gestation (1). PE is a complication of pregnancy constituting a major cause of maternal and fetal morbidity and mortality. Several etiologies have been implicated in the development of preeclampsia including abnormal trophoblast invasion of uterine blood vessels and immunological intolerance between fetoplacental and maternal tissues (2).

Endothelial cell dysfunction and inflammation are considered to have a role in the pathophysiology of PE (3,4). A generalized activation of circulating leukocytes, characteristic of inflammation, has been found during PE. Moreover, increased concentrations of CRP and inflammatory cytokines have been reported in PE women (5).

CRP is produced by the liver and the production is stimulated by the inflammatory cytokines interleukin-6 and TNF-alpha. CRP is a sensitive marker of inflammatory activity in the body. CRP level increases during inflammatory response to tissue injury or infection (6).

The aim of this study was to determine serum CRP level in PE, and its association with the severity of the disease and with biochemical and clinical parameters.

Patients and Methods

This cross-sectional study was conducted in Afzalipour an University Hospital, Kerman, Iran, over a 1-year period from March 2006 to March 2007. The study was performed in three groups of women. Group1 consisted of 43 women with mild preeclampsia at the time of admission; group 2 included 43 women with severe preeclampsia, and group 3 consisted of 43 healthy normotensive women in the third trimester of their pregnancy.

*Corresponding Author: Fatemeh Mirzaie

Department of Obstetric and Gynecology, School of Medicine, Kerman University of Medical Sciences, Kerman, Iran
Tel: +98 341 2447874, Fax: +98 341 3222763, E-mail: mirzaie_fatemeh@yahoo.com

Maternal serum C- reactive protein levels with severity of preeclampsia

Normal pregnancy was diagnosed on the basis of clinical and biochemical findings.

All groups had gestational ages of 28-40 weeks, not in labor and singleton. Patients with a history of diabetes, renal disease, chronic hypertension, systemic lupus erythematosus, systemic infection and cardiovascular disease were excluded during routine interviews, clinical investigations and laboratory tests.

PE was diagnosed according to the following criteria: a blood pressure higher than 140/90 mm Hg and proteinuria more than 300 mg/24h or protein dipstick $\geq 1+$ on ≥ 2 midstream samples 6 hours apart. Severe PE was classified if diastolic blood pressure increased to at least 110mmHg, Proteinuria > 2000 g/24h or protein dipstick $\geq 2+$ on ≥ 2 midstream samples 6 hours apart, and the presence of headache, visual disturbances, epigastric pain, oliguria (30cc/l), increased bilirubin, elevated serum creatinine levels (≥ 0.9 mg/dl), thrombocytopenia (< 100.000 mm³), and elevated aspartate and alanine transaminase levels.

Blood samples were drawn on admission in the morning after a 6h fasting and before magnesium sulfate administration. Serum CRP levels were measured by high CRP sensitivity kits using an immunoturbidimetric method with a detection limit of 0.03 mg/l.

The results were expressed as mean \pm SD and analyzed by an independent samples *t*-test. The correlation analysis was done by spearman's test.

All statistical analyses were carried out by using a SPSS software, version 13. The level of significance was set at $P < 0.05$.

Results

The demographic and clinical parameters of the groups are summarized in table 1.

Maternal age was not significantly different between the groups ($P > 0.05$, table 1). Mean systolic and diastolic blood pressures were higher in PE groups as compared to healthy women values ($P = 0.001$, Table 1).

Gestational age was significantly different between the groups ($P = 0.0001$, Table 1).

Serum CRP levels in mild and severe PE were markedly higher than those of normal third trimester pregnant women (Table 2).

The hemoglobin levels, platelet count and blood urea nitrogen, creatinine, aspartate transaminase (AST), alanine transaminase (ATT), and Alkaline phosphatase were higher in mild and severe PE (Table 2). Bilirubin levels were found to be similar in three groups.

Table 1. Demographic and clinical parameters of groups

	Healthy group	Mild PE	Sever PE	P-Value
Maternal age	26.31 \pm 5.13	27.6 \pm 6.14	28.62 \pm 5.40	NS
Gestational age	38.63 \pm 1.87	36.86 \pm 3.45	35.34 \pm 3.66	$P = 0.0001$
Systolic BP (mmHg)	109.17 \pm 9.81	137.38 \pm 4.96	162.50 \pm 14.15	$P = 0.001$
Diastolic BP(mmHg)	71.90 \pm 7.56	88.21 \pm 8.17	96.43 \pm 1.42	$P = 0.001$
Birth weight (kg)	3.16 \pm 0.53	2.89 \pm 0.53	2.64 \pm 0.7	$P = 0.0001$

NS: Non Significant, BP: Blood Pressure

Table 2. Biochemical parameters of study groups

	Healthy group (n=43)	Mild preeclampsia (n=43)	Sever preeclampsia (n=43)	P-Value
CRP(mg/l)	3.42 \pm 5.48	14.28 \pm 11.62	34 \pm 25.27	$P = 0.003$
ALT(u/l)	20.79 \pm 5.93	21.26 \pm 5.69	37.33 \pm 31.39	$P = 0.001$
AST(u/l)	15.93 \pm 6.63	17.88 \pm 5.23	38.05 \pm 38.77	$P = 0.0001$
Creatinine (mg/dl)	0.75 \pm 0.24	0.79 \pm 0.20	0.93 \pm 0.25	$P = 0.002$
Blood urea nitrogen (mg/dl)	22.88 \pm 7.54	23.33 \pm 7.77	28.50 \pm 8.59	$P = 0.002$
Alkaline phosphatase	253.43 \pm 84.50	242.12 \pm 113.33	308.48 \pm 142.49	$P = 0.007$
Hemoglobin (mg/dl)	12.59 \pm 7.54	13.47 \pm 1.35	15.03 \pm 1.35	$P = 0.0001$
Total Bilirubin (mg/dl)	0.41 \pm 0.12	0.4 \pm 0.2	0.46 \pm 0.20	$P = 0.19$

AST: Aspartate Transaminase, ALT: Alanine Transaminase

In mild to severe PE groups, correlation tests showed a positive correlation between serum CRP levels and systolic blood pressure ($r = 0.6, p = 0$), diastolic blood pressure ($r = 0.5, P = 0$), creatinine levels ($r = 0.2, p = 0.003$), blood urea nitrogen ($r=0.2, p=0.004$), urinary protein excretion ($r = 0.5, P = 0.00$), ALT ($r = 0.2, P = 0.006$), AST ($r = 0.3, P = 0/001$), Hemoglobin level ($r = 0.2, P = 0.001$), and Alkaline phosphatase ($r = 0.19, r = 0.003$). There was a negative correlation between CRP and birth weight ($r = -0.09, P = 0.01$) and gestational age ($r = -0.4, P = 0$).

Discussion

Preeclampsia is a disease of pregnancy associated with endothelial cell damage. There is increasing evidence that preeclampsia is a systemic inflammatory disease (2).

Studies have shown that markers of endothelial activation or inflammation have an active role in preeclampsia (7).

CRP, a sensitive marker of tissue damage and inflammation, was proposed to play a role in eliciting the inflammatory response characteristics of preeclampsia (2).

In this study, we have clearly shown that serum CRP levels were higher in Preeclampsia woman as compared to healthy pregnant controls. Similarly, there are studies reporting that serum CRP levels increase in preeclamptic woman (1, 2). There are few studies concerning correlation of CRP levels with severity of preeclampsia. Determination of serum CRP levels in the third trimester pregnant women proved of great value in predicting the prognosis of preeclampsia. Proteinuria and blood pressures are used as parameters for severity of PE (8).

We found a positive correlation between serum levels of CRP and diastolic and systolic blood pressure and urinary protein excretion in PE. Endothelial dysfunction of PE has been associated with an exaggerated maternal inflammatory response to pregnancy. It has been hypothesized that placental hypoxia, resulting from uteroplacental arterial insufficiency, amplifies the release of inflammatory stimuli into the maternal circulation (9).

In our study, there was a positive correlation between CRP levels and biochemical parameters (ALT, AST, Bun-creatinine, hemoglobin, proteinuria)

Kumru *et al.* (2006) (1) showed a positive correlation between serum CRP levels and diastolic blood pressure and urinary protein excretion (1). Another study found a significant correlation between mean arterial pressure and CRP ($r = 0.515, P = 0.0001$) in pregnancies compli-

cated with preeclampsia. In this study, correlation of CRP level with other biochemical and clinical parameters of severe preeclampsia was not evaluated (2).

Erren *et al.* (1999) reported that inflammatory profile was more pronounced when the endothelial damage was more advanced. It is well known that renal dysfunction usually occurs in PE, especially in its severe forms. In cases of renal dysfunction, endothelial dysfunction and increased inflammatory markers such as CRP have been shown (10-12).

The mean gestational age was less in our study for the Preeclamptic women than the healthy (healthy = 38.68, mild = 36.86 severe = 35.34, $P = 0.0001$) and in the preeclamptic groups, there was a statistically significant negative correlation between gestational age and maternal CRP concentration ($r = -0.4, P = 0$)

The higher CRP levels in patients with Preeclampsia delivering earlier in pregnancy could probably be a marker of disease severity and a marker of an excessive inflammatory response in patients with the most severe PE disease delivering prematurely compared with patients with less severe disease who deliver later in pregnancy (6).

Our findings of elevated serum levels of CRP are associated with low birth weight and a negative correlation between serum CRP levels and weight of the newborns in preeclampsia groups. It is well known that PE increases the risks of intrauterine growth restriction and low birth weight (13). Kurmus *et al.* (2006) found a negative correlation between serum CRP and weight and length of the newborns in the PE group in comparison with the control group (1). CRP level, a marker of inflammatory response, is raised in healthy pregnant women compared with non-pregnant women. However, factors such as age, labor, infections and medical disease are associated with raised concentrations of CRP. In this study, all groups were matched for maternal age, none had chronic medical disorders or were in labor (14). However, we found an evidence for presence of inflammation in preeclampsia. CRP levels correlated positively with the severity of disease. We also showed a positive correlation between serum CRP and biochemical and clinical parameters in preeclampsia.

References

1. Kumru S, Godekmerdan A, Kutlu S, Ozcan Z. Correlation of maternal serum high-sensitive C-reactive protein levels with biochemical and clinical parameters in preeclampsia. *Eur J Obstet Gynecol Reprod Biol* 2006; 124(2): 164-7.

Maternal serum C- reactive protein levels with severity of preeclampsia

2. Ustun y, Engin-ustun Y, Kamaci M. Association of fibrinogen and C-reactive protein with severity of preeclampsia. *Eur J Obstet Gynecol Reprod Biol* 2005; 121(2): 154-8.
3. Creasy RK, Resnick R, Iams JD, editors. *Maternal-Fetal Medicine: Principles and Practice*. 5th ed. Philadelphia: WB Saunders; 2004.
4. James DK, Steer PJ, Weiner CP, Gonik B, editors. *High Risk Pregnancy Management Options*. 3rd ed. Philadelphia: WB Saunders; 2005.
5. García RG, Celedón J, Sierra-Laguado J, Alarcón MA, Luengas C, Silva F, et al. Raised C-reactive protein and impaired flow-mediated vasodilation precede the development of preeclampsia. *Am J Hypertens* 2007; 20(1): 98-103.
6. Braekke K, Holthe MR, Harsem NK, Fagerhol MK, Staff AC. Calprotectin, a marker of inflammation, is elevated in the maternal but not in the fetal circulation in preeclampsia. *Am J Obstet Gynecol* 2005; 193(1): 227-33.
7. Teran E, Escudero C, Moya W, Flores M, Vallance P, Lopez-Jaramillo P. Elevated C-reactive protein and pro-inflammatory cytokines in Andean women with preeclampsia. *Int J Gynaecol Obstet* 2001; 75(3): 243-9.
8. ACOG Committee on Obstetric Practice. ACOG practice bulletin. Diagnosis and management of preeclampsia and eclampsia. Number 33, January 2002. American College of Obstetricians and Gynecologists. *Int J Gynaecol Obstet* 2002; 77(1): 67-75.
- 9 Redman CW, Sacks GP, Sargent IL. Preeclampsia: an excessive maternal inflammatory response to pregnancy. *Am J Obstet Gynecol* 1999; 180(2 Pt 1): 499-506.
10. Erren M, Reinecke H, Junker R, Fobker M, Schulte H, Schurek JO, et al. Systemic inflammatory parameters in patients with atherosclerosis of the coronary and peripheral arteries. *Arterioscler Thromb Vasc Biol* 1999; 19(10): 2355-63.
11. Stam F, van Guldener C, Schalkwijk CG, ter Wee PM, Donker AJ, Stehouwer CD. Impaired renal function is associated with markers of endothelial dysfunction and increased inflammatory activity. *Nephrol Dial Transplant* 2003; 18(5): 892-8.
12. Stuveling EM, Hillege HL, Bakker SJ, Gans RO, De Jong PE, De Zeeuw D. C-reactive protein is associated with renal function abnormalities in a non-diabetic population. *Kidney Int* 2003; 63(2): 654-61.
13. Naicker T, Khedun SM, Moodley J, Pijnenborg R. Quantitative analysis of trophoblast invasion in preeclampsia. *Acta Obstet Gynecol Scand* 2003; 82(8): 722-9.
14. Hwang HS, Kwon JY, Kim MA, Park YW, Kim YH. Maternal serum highly sensitive C-reactive protein in normal pregnancy and pre-eclampsia. *Int J Gynaecol Obstet* 2007; 98(2): 105-9.