

THE INVESTIGATION OF CRITERION VALIDITY OF THE MODIFIED ASHWORTH SCALE USING THE ALPHA MOTONEURON EXCITABILITY INDICATORS

N. Ghotbi¹, M. R. Hadian^{1*}, G. R. Olyaei¹, H. Bagheri¹, S. Talebian¹, N. Nakhostin-Ansari¹ and S. Nafissi²

1) Electrophysiology Laboratory, Faculty of Rehabilitation, Medical Sciences/University of Tehran, Tehran, Iran

2) Department of Neurology, School of Medicine, Medical Sciences/University of Tehran, Tehran, Iran

Abstract- Despite the large body literature focused on the modified Ashworth Scale (MAS), validity of the scale for measuring spasticity is still questionable. The purpose of this study was to investigate the validity of the MAS by using both the conventional (Hmax/Mmax ratio) and the new indicators (Hslp/Mslp) of motoneuron excitability. Main outcome measures were the MAS and electrophysiologic assessments. The latter was performed using both conventional and new indicators of alpha motoneuron excitability. Data on 20 hemiplegic patients (17 men and three women) were analyzed. Correlation between the MAS and either soleus Hmax/Mmax or soleus Hslope/Mslope was not significant. In 10 patients whose H-reflex could be evoked bilaterally, only new spinal excitability indicator showed significant difference between the affected and non-affected sides. Based on the results of this study, there is no relationship between the MAS and the indicators of alpha motoneuron excitability. This research suggests that the MAS is not a valid measure for the assessment of spasticity in ankle plantar flexors. *Acta Medica Iranica* 2007; 45(4): 290-294.

© 2007 Tehran University of Medical Sciences. All rights reserved.

Key words: Stroke, Spasticity, Ashworth Scale, Hoffmann Reflex, Motoneuron excitability

INTRODUCTION

Stroke is a common disorder, which causes long-term disability among adults (1). One of the positive features of stroke is spasticity. Spasticity has been defined as a “motor disorder characterized by a velocity-dependent increase in tonic stretch reflexes (muscle tone) with exaggerated tendon jerks resulting from hyper excitability of the stretch reflex as one component of the upper motor neuron (UMN) syndrome (2). Modified Ashworth scale (MAS) is a clinical rating test which is based on the assessment of resistance to passive movement at a joint.

Currently, it is the most widely used test for measurement of muscle spasticity in research and clinical practice (3-8). However, despite the large body literature focused on the MAS, validity of the scale for measuring spasticity is still questionable (3, 6). Since spasticity is associated with increased excitability of alpha motoneurons (3) and H-reflex indirectly measures the spinal excitability; some researches mainly used the conventional indicator of H-reflex (Hmax/Mmax ratio) for assessing the spasticity (3, 9). However, as previously reported, Hmax/Mmax ratio may lead to false assessments of motor neuron pool excitability (10).

In order to overcome this obstacle, new indicator of Hoffmann reflex (*i.e.* Hslope/Mslope ratio) has been introduced. It is reported that this parameter is a more accurate and sensitive measure of spinal excitability (10-12).

Received: 2 Dec. 2005, Revised: 14 Jul. 2006, Accepted: 4 Sep. 2006

*** Corresponding Author:**

Mohammad Reza Hadian, Electrophysiology Laboratory, Faculty of Rehabilitation, Medical Sciences/Tehran University, Tehran, Iran
Tel: +98 21 77536134
Fax: +98 21 77534133
E-mail: hadianrs@sina.tums.ac.ir

In this respect, the main objective of current research was to investigate the validity of this scale by using both the conventional and new indicators of soleus Hoffmann reflex.

MATERIALS AND METHODS

Twenty hemiplegic patients due to first stroke participated in this correlational study. This study was approved by the Ethical Committee of Tehran University and each subject provided informed written consent. Inclusion criteria: absence of ankle fix contracture, no use of drugs which affects neuromuscular system and being in spastic phase.

Procedures

All information with regard to gender, weight, height and post stroke duration was recorded. Thereafter, clinical and electrophysiological assessments were performed in the same session and under the same conditions, respectively.

Clinical assessment

Patients were lying flat in supine position with socks and shoes removed. As the level of muscle tone can be affected by the volume of bladder, patients were asked to empty their bladder.

Spasticity of the calf muscles was rated according to the MAS. This scale measures the resistance to passive movement and is a 6-point scale from 0 (no increase in muscle tone) to 4 (affected part[s] rigid in flexion or extension) (13).

All tests were carried out by one investigator to eliminate inter-rater variability (3).

Electrophysiological assessment

Patients were comfortably positioned in prone lying and the feet suspended over the end of the bed. The skin temperature over the calf muscles was monitored via a digital thermometer. Paired surface electrodes (silver-silver chloride) were placed to the belly of the soleus muscle according to Hugon classic protocol (Hugon, 1973). Tibial nerve in the popliteal fossa was stimulated via a cathode placed on the skin overlying the nerve (with a rectangular electric pulse of 0.5 ms duration and a stimulus frequency of 1 per five seconds). The anodal electrode was placed on the

quadriceps muscle above the patella. A cuff type ground electrode was placed between the stimulating cathode and the EMG electrodes.

The stimulus strength was gradually increased from below threshold to supramaximal. Three pulses were delivered sequentially at each stimulus intensity, and the mean was calculated to allow for normal variation in H-reflex amplitude at rest.

EMG signals were filtered (5, 5000Hz), and amplified (gain 500-2500 for H-reflex) with Toennies (Neuroscreen Plus, Germany) and stored for further analysis.

Data Analysis

Peak-to-peak H-reflex and M-wave amplitudes were measured from the largest positive and negative deflection from the baseline. The mean of the three H-reflexes collected at each stimulus intensity was expressed as a percentage of the maximal M-wave (Mmax). Stimulus intensity was expressed relative to the threshold intensity for an M-wave (Mth) (12, 10). For each patient, the data points being in the middle of the linear portion of the H-reflex recruitment curve were analyzed by linear regression. In this way, the rate of the ascending slope rise was calculated.

The rate of rise of the H-reflex recruitment curve described the increase in H-reflex amplitude per unit of threshold stimulus intensity and was called H-slope (Hslp) (Fig. 1).

The rate of rise of the linear portion of the M-wave recruitment curve was calculated in a similar manner and called M-slope (Mslp) (Fig. 1).




Fig. 1. Recruitment curve in a patient.

Validity of modified Ashworth scale

Figure 1 shows the recruitment curves of H-reflex and M-response in the affected side of a patient who had an MAS score of 2. The two lines represent the slope of H-curve and M-curve (H-slope a M-slope).

Statistical test

To determine the criterion validity of the scale, correlation between it and each of the alpha motoneuron excitability measures was calculated by Spearman rank correlation test.

Analysis of differences between affected and unaffected sides was made with the paired *t* test. Data were analyzed using the SPSS statistical package (version 11.5).

RESULTS

Twenty hemiplegic patients (3 women, 17 men) were investigated. Twelve patients had right hemiplegia, and eight subjects had left hemiplegia. Mean age,

height and weight were 61.80 ± 11.98 years, 168.79 ± 11.38 cm and 75.89 ± 12.86 kg, respectively.

Descriptive information about the characteristics of the study population is included in table 1. Range of the MAS scores was 0 to 3. Mean \pm SD values of soleus Hmax/Mmax and Hslope/Mslope ratios were 0.45 ± 0.25 and 1.56 ± 1.45 , respectively. There was no significant correlation between degree of spasticity, as rated by the MAS and either Hmax/Mmax ratio or Hslope/ Mslope ratio ($P > 0.05$).

In 10 of 20 patients, H-reflex could be evoked bilaterally. Thus, mean \pm SD values of the soleus Hmax/Mmax ratio (unaffected side: 0.25 ± 0.23 , affected side: 0.38 ± 0.19) and Hslope/Mslope ratio (unaffected side: 0.46 ± 0.30 , affected side: 1.05 ± 0.53) was calculated in ten patients. Only Hslope/Mslope ratio showed a significant difference between two sides ($t = 4.2$, $df = 9$, $P = 0.002$) (Fig. 2).

Table 1. Descriptive data in hemiplegic patients

Patient	Age	Stroke Duration	MAS scores	Hmax/Mmax	Hslope/Mslope
1	47	4	1 ⁺	0.60	1.60
2	54	22	2	0.26	0.31
3	69	12	2	0.28	0.59
4	57	29	1	0.63	1.87
5	59	10	2	0.47	1.49
6	64	60	3	0.14	0.24
7	39	26	3	0.91	4.69
8	72	1	1	0.64	2.51
9	81	54	3	0.49	1.36
10	70	8	1 ⁺	0.32	2.01
11	53	13	0	0.04	0.05
12	42	12	1	0.45	0.83
13	68	32	1	0.41	1.10
14	47	1	1 ⁺	0.23	0.85
15	80	12	1	0.09	0.36
16	76	9	0	0.31	1.84
17	67	36	2	0.78	1.48
18	65	60	1 ⁺	0.46	1.31
19	63	5	0	0.68	2.46
20	63	9	2	0.82	5.84
Total*	61.80 ± 11.98 (39-81)	20.75 ± 18.91 (1-60)	(0-3)	0.45 ± 0.25 (0.04-0.91)	1.56 ± 1.45 (0.07-5.84)

Abbreviations: Hslp, H slope; Mslp, M slope.

* Data for total are given as mean \pm SD (range).




Fig. 2. Mean \pm SD values of conventional (A) and new (B) indicators of alpha motoneuron excitability in the affected and unaffected of 10 patients. Number 1 and 2 indicates the values for affected and unaffected sides, respectively.

DISCUSSION

Up to the present, this research was the first to use recruitment curves for determining the validity of the MAS. Previous studies, investigated the relationship between MAS and either biomechanical measures (6) or traditional parameter of H-reflex (3, 7, 9, 14). Most of which showed low to moderate correlation (6, 14, 9) and some presented no correlation (7, 9).

Our findings showed that there was no significant correlation between the MAS scores and either Hmax/Mmax ratio or Hslope/Mslope ratio. New indicator of spinal excitability was calculated via recruitment curves of H-reflex and M-wave and linear regression line of them. Although, the Hslope/Mslope ratio is more sensitive than Hmax/Mmax ratio for the assessment of motoneuron excitability, the results of the present study showed that the Hslope/Mslope ratio has no relationship with

the MAS scores. Since the MAS relies on subjective assessment of reflex activity, lack of its correlation with neurophysiologic tests may relate to problems inherent to the scale itself.

Only Hslope/Mslope ratios showed significant difference between the affected and unaffected sides in our patients. This finding was in agreement with Higashi et.al study (12) and may suggest that the rate of excitability of motoneurons is not symmetrical in both sides. The increase of spinal excitability in the affected side could be related to a decreased inhibitory control on the lower motoneuron or to an increase of stretch reflex excitability secondary to a presynaptic inhibition (15).

In the present study, in order to increase the reliability of experiments, the following points were considered; both tests were carried out in the same session and under the same conditions and the factors which could influence the H-reflex test (e.g. head and limb position) (16) and the MAS test (e.g. bladder status) were controlled. Thus, in conclusion, no correlation between the scale and spinal excitability measures suggests that MAS does not provide a valid measure for evaluating spasticity in ankle plantar flexors.

Acknowledgment

This project was supported by a grant from the Postgraduate Studies and Research Program, Medical Sciences/University of Tehran, Tehran, Iran. The authors would like to acknowledge the generous assistance of the staff of Faculty of Rehabilitation, Tehran University.

Conflict of interests

We have no conflict of interests.

REFERENCES

1. Hallett M, Wassermann EM, Cohen LG, Chmielowska J, Gerloff C. Cortical mechanisms of recovery of function after stroke. *Neurorehabilitation*. 1998; 10:131-142.
2. Lance JW. Symposium synopsis. In: Feldman RG, Young RR, Koala WP, eds. *Spasticity: disordered motor control*. Chicago: Year Book Medical; 1980. P. 487- 489.

Validity of modified Ashworth scale

3. Bakheit AM, Maynard VA, Curnow J, Hudson N, Kodapala S. The relation between Ashworth scale scores and the excitability of the alpha motor neurones in patients with post-stroke muscle spasticity. *J Neurol Neurosurg Psychiatry*. 2003 May; 74(5):646-648.
4. Sehgal N, McGuire JR. Beyond Ashworth. Electrophysiologic quantification of spasticity. *Phys Med Rehabil Clin N Am*. 1998 Nov; 9(4):949-479.
5. Pandyan AD, Price CI, Rodgers H, Barnes MP, Johnson GR. Biomechanical examination of a commonly used measure of spasticity. *Clin Biomech (Bristol, Avon)*. 2001 Dec; 16(10):859-865.
6. Pandyan AD, Price CI, Barnes MP, Johnson GR. A biomechanical investigation into the validity of the modified Ashworth Scale as a measure of elbow spasticity. *Clin Rehabil*. 2003 May; 17(3):290-293.
7. Pisano F, Miscio G, Del Conte C, Pianca D, Candeloro E, Colombo R. Quantitative measures of spasticity in post-stroke patients. *Clin Neurophysiol*. 2000 Jun; 111(6):1015-1022.
8. Johnson GR. Measurement of spasticity. In: Barnes MP, Johnson GR, eds. *Upper motor neuron syndrome and spasticity: clinical measurement and neurophysiology*. Cambridge: Cambridge University Press; 2001. P. 79-95.
9. Pizzi A, Carlucci G, Falsini C, Verdesca S, Grippo A. Evaluation of upper-limb spasticity after stroke: A clinical and neurophysiologic study. *Arch Phys Med Rehabil*. 2005 Mar; 86(3):410-415.
10. Higashi T, Funase K, Kusano K, Tabira T, Harada N, Sakakibara A, Yoshimura T. Motoneuron pool excitability of hemiplegic patients: assessing recovery stages by using H-reflex and M response. *Arch Phys Med Rehabil*. 2001 Nov; 82(11):1604-1610.
11. Funase K, Higashi T, Yoshimura T, Imanaka K, Nishihira Y. Evident difference in the excitability of the motoneuron pool between normal subjects and patients with spasticity assessed by a new method using H-reflex and M-response. *Neurosci Lett*. 1996 Jan 19;203(2):127-130.
12. Bradnam L, Rochester L, Vujnovich A. Manual cervical traction reduces alpha-motoneuron excitability in normal subjects. *Electromyogr Clin Neurophysiol*. 2000 Jul-Aug; 40(5):259-266.
13. Bohannon RW, Smith MB. Interrater reliability of a modified Ashworth scale of muscle spasticity. *Phys Ther*. 1987 Feb; 67(2):206-207.
14. Allison SC, Abraham LD. Correlation of quantitative measures with the modified Ashworth scale in the assessment of plantar flexor spasticity in patients with traumatic brain injury. *J Neurol*. 1995 Oct; 242(10):699-706.
15. Schieppati M. The Hoffmann reflex: a means of assessing spinal reflex excitability and its descending control in man. *Prog Neurobiol*. 1987;28(4):345-376.
16. Voerman GE, Gregoric M, Hermens HJ. Neurophysiological methods for the assessment of spasticity: the Hoffmann reflex, the tendon reflex, and the stretch reflex. *Disabil Rehabil*. 2005 Jan 7-21;27(1-2):33-68.