

Effects of Different Doses of Hyaluronan on Human Sperm Motility, Vitality and Morphology

Amaneh Mohammadi Roushandeh¹, Farnaz Sohrabvand², Saeid Amanpour², Parichehr Pasbakhsh³,
Abdollah Sobhani^{4*}, Hossein Nikzad⁵, and Aligholi Sobhani⁶

¹ Department of Anatomy, School of Medicine, Tabriz University of Medical Sciences, Tabriz, Iran

² VALI-e-ASR Reproductive Health Research Center, Tehran University of Medical Sciences, Tehran, Iran

³ Department of Anatomy, Faculty of Medicine, Tehran University of Medical Sciences, Tehran, Iran

⁴ Department of Animal Sciences, Faculty of Agriculture, University of Mohaghegh Ardabili, Ardabil, Iran

⁵ Department of Anatomy, Faculty of Medicine, Kashan University of Medical Sciences, Kashan, Iran

⁶ Department of Anatomy and VALI-e-ASR Reproductive Health Research Center, Faculty of Medicine, Tehran University of Medical Sciences, Tehran, Iran

Received: 12 Mar. 2008; Received in revised form: 4 Apr. 2008; Accepted: 14 May 2008

Abstract- Important aspect of sperm function such as motility and capacitation appear to be mediated at least partially through hyaluronic acid (HA). Present study investigated effects of different doses of HA on sperm motility and vitality in human. Sperm was obtained from 20 male from IVF clinic in Imam Khomeini Hospital. Sperm motility and vitality in human semen was analyzed according to WHO criteria before and 4 hours after treatment with different doses of HA (0.750, 1000 and 1250 µg/ml). The results showed that in 1000 µg/ml the percent of stage 3 and 4 increased compare to control group. Percent of stage 1 and 2 decreased in group with 1000 µg/ml HA, there was an increase in the percentage of stage 3 and 4 and decrease in percentage of stage 1 and 2 compare to control. In the group treated with 1250 µg/ml stage 1 and 2 increased while stage 3 and 4 decreased. Vitality in all groups decreased except of the group treated with 1000 µg/ml HA. The group with 1250 µg/ml showed significantly decrease in vitality compare to fresh group ($P < 0.05$). The present study showed that the effects of HA on sperm motility and vitality is dose dependant and 1000 µg/ml HA had the effective role on sperm parameters.

© 2009 Tehran University of Medical Sciences. All rights reserved.

Acta Medica Iranica 2009; 47(5): 369-372.

Key words: Human sperm, hyaluronan, sperm parameters

Introduction

Glycosaminoglycans (GAGs) are essential components of the extracellular matrix, contributing to cell recognition, cellular adhesion and growth regulation. The GAGs can be classified by their sugar residues and other characteristics such as sulfation into four main groups of GAGs including the non-sulfated glycosaminoglycan, hyaluronic acid (HA), or hyaluronan (1). Important aspects of sperm function such as motility (2, 3) and capacitation (4, 5) appear to be mediated, at least partially, through HA. Addition of HA to fresh (6,7) and frozen-thawed (8) sperm improves sperm motility in medium. Interestingly, HA not only improves sperm motility after freezing and thawing procedures, but it further appears

to stabilize spermatozoa with already declining motility (7). The enhancement of sperm motility and velocity occurred as a direct response to HA. There is an instantaneous increase in sperm velocity and tail cross-beat frequency upon HA exposure and when, after density gradient centrifugation, the HA-exposed sperm is transferred to a standard medium, the motility and velocity properties return to those of the control sperm. It can be concluded that HA effects on sperm are likely to be receptor mediated, in line with the evidence established by various laboratories for the presence of the HA receptor in human sperm (8). Concurrently, with the sperm maturation studies, in another line of experiments, scientists have investigated the effects of HA on human sperm function. Exogenous HA decreased polyspermy during

*Corresponding Author: Aligholi Sobhani

Department of Anatomy and VALI-e-ASR Reproductive Health Research Center, School of Medicine, Tehran University of Medical Sciences, Tehran, Iran

Tel: +98 21 64432348, Fax: +98 21 64432348, E-mail: sobhania@tums.ac.ir

Hyaluronan and sperm parameters

conventional porcine IVF (4), an effect probably related to the capacity of HA to delay sperm capacitation and/or the acrosome reaction (4, 5). It has been reported that effects of hyaluronan on sperm is dose dependant (9). There isn't any report about the optimal dose of hyaluronan on human sperm motility, viability and morphology. The aim of the present study was to test the hypothesis that supplementing with different dose of HA has an important role on the viability and motility of sperm.

Patients and Methods

This study was initially approved in the ethical and scientific committee of the Vali-e-Asr Reproductive Health Research Center, Tehran, Iran. Semen samples were obtained from 20 couples referred to Vali-e-Asr Reproductive Health Research Center for ICSI and IVF treatment. The semen samples were collected by masturbation after 3-4 days of abstinence on the day of oocyte recovery. Routine semen analysis was carried out by light microscopy according to World Health Organization (WHO) criteria. After ART treatment, the remaining samples were used for this research. Each sample was divided into five groups. The Group 1 (fresh control): These samples were analyzed freshly without any treatment. The Group 2 (control): They were maintained in HTF media (Sigma) without any hyaluronan (Sigma) supplementation for 2 h. In groups 3, 4 and 5, HA with the doses of 750, 1000 and 1250 µg/ml were added into HTF media, respectively. After 4 h incubation, the sperms were assessed for motility, vitality and morphology.

Sperm analysis

Sperm motility was analyzed according to WHO manual. For evaluating sperm viability, we used the eosin staining. Sperm and eosin (Sigma) were mixed (3:1) and number of sperm was counted by 400 magnifica-

tions by an invert microscope rapidly. Sperm smear was prepared on the slide for morphology analyzing. After fixing with 1:1 volume of alcohol (sigma) and ether (sigma), the slides were stained with papanicolaou methods. Numbers of coiled sperm, stumped tail sperm, cytoplasmic droplet and remnant were analyzed by binocular microscope with 400 magnifications.

Hyaluronan treatment

After evaluation of fresh sperm, the samples were treated with different doses of Hyaluronan (0, 750, 1000 and 1250µg/ml). The samples were put in incubator (37c° and 5% Co₂ for 4 hours). After 4 hours the different groups of sperm were analyzed again.

Statistical analysis

Data were analyzed by One-Way ANOVA followed up with post hoc test (Tukey). Statistical significance was set at $P < 0.05$.

Results

Sperm motility

Table 1 indicates the various motility stages of sperm in different groups. As shown in this table, percentage of stage 1 in group 4 (1000 µg/ml) was decreased significantly compared to control groups ($P < 0.05$) but in group 5 (1250 µg/ml) the stage 1 significantly increased. The percentage of stage 2 decreased in group 4 compared to other groups. In group 4, stage 4 motility significantly increased compared to all off other groups. In group 5 there was significant decrease in stages 4 motility ($P < 0.05$).

Sperm viability

As shown in figure 1, viability in all groups after treatment with HA was decreased except for group 4. In group 5 significant decrease ($P < 0.05$) in percentage of viability compared to fresh group was observed.

Table 1. Various stages of motility in different groups

Groups	Fresh control	Control after 4 hr	750 µg/ml HA	1000 µg/ml HA	1250 µg/ml HA
Stages					
1	14%	15%	16%	12%	21%
2	16%	17%	17%	10%	25%
3	25%	28%	29%	33%	28%
4	45%*	40%	38%	45%	26%




Figure 1. Percent of viability in different groups

Discussion

Hyaluronan (HA) is a linear polysaccharide which is a mediator of cell locomotion (9, 10). HA also occurs physiologically in the reproductive system in cervical mucus, the cumulus and the follicular and seminal fluids. The effects of HA on human spermatozoa have been explored in other clinical and research applications (12-15). There are some reports regarding the effect of different doses of HA on sperm function(8,9). Sbracia *et al.* (8) showed that addition of 250µg/ml HA provided a significant motility increase for at least 4 h after the Percoll purification of spermatozoa. However, what remains to be elucidated is determining the optimum doses of HA on sperm function. Therefore, we used different doses of HA were applied on sperm to evaluate their effect on motility, viability and morphology. The results of this study showed that effect of HA on sperm motility were dose dependant and 1000 µg/ml was optimal dose to improve the fast motility (stages 3 and 4). Furthermore, results showed that 1250µg/ml had a toxic effect on sperm and motility was decreased significantly. The mechanisms behind a HA-induced increase in sperm motility remains to be determined. A HA-receptor has been localized along the tail, mid piece, and head of human spermatozoa (16). Blocking this receptor with mono-specific antibodies resulted in inhibition of sperm motility. Moreover, enhanced phosphorylation of the 34 kDa HA-binding protein (HABP) has been observed in response to HA supplementation, suggesting an important role of this glycosaminoglycan in initiating the transduction of signals controlling important processes of sperm physiology, such as motility (17, 18) that may result in an increase of the ATP levels and improved flagellar function(8). Bakhtiari *et al.* showed that, the hyaluronan may have an influence on motility, viability

and fertilizing rate of mouse sperm and the dose of 750 µg/ml had a significant effect on these factors. The results of this study, in addition, showed that viability in hyaluronan group after four hours decreased except for 1000 µg/ml. Sbracia *et al.* (8) showed that viability of incubated with HA increased. These results are in agreement with those of other research (8, 9). This effect could be related to protective role of HA on sperm membrane. According to our study it seems that 1000 µg/ml is the optimal dose that affects motility and vitality of human sperm. In this investigation different doses didn't have an effect on sperm morphology. Previous studies have shown that sperm morphology is controlled by spermiogenic events such as acrosome formation and chromatin rearrangement. Protamine deficiency and aberrant P1/P2 ratios are negatively related to abnormal morphology (19). After ejaculation sperm morphology isn't affected by HA but it is possibly affected by osmolarity of sperm medium preparation. It seems that HA has a substantial advantage over other sperm motility enhancers, which have a temporary effect. Unlike other motility enhancers, some of which cause short-term maintenance of improvement after removal of the agent, HA preserves motility in spermatozoa and therefore it can be useful in assisted reproduction technique (ART) in utilizing, especially when thawed semen samples are used. Also, when preparing sperm for IUI, IVF, application of HA which leads to extended motility will increase fertilizing efficiency, while in intracytoplasmic sperm injection (ICSI) the improved sperm motility will enhance the selection of viable spermatozoa.

Acknowledgements

This project financially supported by research deputy of Tehran University of Medical Sciences. So, we would like to thank for their helps.

References

1. Erlinger R. Glycosaminoglycans in porcine lung: an ultrastructural study using cupromeronic blue. *Cell Tissue Res* 1995;281(3):473-83.
2. Vines CA, Li MW, Deng X, Yudin AI, Cherr GN, Overstreet JW. Identification of a hyaluronic acid (HA) binding domain in the PH-20 protein that may function in cell signaling. *Mol Reprod Dev* 2001;60(4):542-52.
3. Ghosh I, Bharadwaj A, Datta K. Reduction in the level of hyaluronan binding protein 1 (HABP1) is associated with loss of sperm motility. *J Reprod Immunol* 2002;53(1-2):45-54.

Hyaluronan and sperm parameters

4. Suzuki K, Asano A, Eriksson B, Niwa K, Nagai T, Rodriguez-Martinez H. Capacitation status and in vitro fertility of boar spermatozoa: effects of seminal plasma, cumulus-oocyte-complexes-conditioned medium and hyaluronan. *Int J Androl* 2002;25(2):84-93.
5. Suzuki K, Eriksson B, Shimizu H, Nagai T, Rodriguez-Martinez H. Effect of hyaluronan on monospermic penetration of porcine oocytes fertilized in vitro. *Int J Androl* 2000;23(1):13-21.
6. Huszar G, Willetts M, Corrales M. Hyaluronic acid (Sperm Select) improves retention of sperm motility and velocity in normospermic and oligospermic specimens. *Fertil Steril* 1990;54(6):1127-34.
7. Bakhtirai M, Sobhani A, Akbari M, Pasbakhsh P, Abbasi A, Hedaiatpour A, et al. The effect of hyaluronic acid on motility, vitality and fertilization capability of mouse sperms after cryopreservation. *Iranian J Reproductive Med* 2007;5(2):45-50.
8. Sbracia M, Grasso J, Sayme N, Stronk J, Huszar G. Hyaluronic acid substantially increases the retention of motility in cryopreserved/thawed human spermatozoa. *Hum Reprod* 1997;12(9):1949-54.
9. Peña FJ, Johannisson A, Wallgren M, Rodriguez-Martinez H. Effect of hyaluronan supplementation on boar sperm motility and membrane lipid architecture status after cryopreservation. *Theriogenology* 2004;61(1):63-70.
10. Turley EA, Austen L, Vandeligt K, Clary C. Hyaluronan and a cell-associated hyaluronan binding protein regulate the locomotion of ras-transformed cells. *J Cell Biol* 1991;112(5):1041-7.
11. Turley EA. Hyaluronan and cell locomotion. *Cancer Metastasis Rev* 1992;11(1):21-30.
12. Karlström PO, Bakos O, Bergh T, Lundkvist O. Intrauterine insemination and comparison of two methods of sperm preparation. *Hum Reprod* 1991;6(3):390-5.
13. Mortimer D, Mortimer ST, Shu MA, Swart R. A simplified approach to sperm-cervical mucus interaction testing using a hyaluronate migration test. *Hum Reprod* 1990;5(7):835-41.
14. Slotte H, Akerlöf E, Pousette A. Separation of human spermatozoa with hyaluronic acid induces, and Percoll inhibits, the acrosome reaction. *Int J Androl* 1993;16(6):349-54.
15. Neuwinger J, Cooper TG, Knuth UA, Nieschlag E. Hyaluronic acid as a medium for human sperm migration tests. *Hum Reprod* 1991;6(3):396-400.
16. Kornovski BS, McCoshen J, Kredentser J, Turley E. The regulation of sperm motility by a novel hyaluronan receptor. *Fertil Steril* 1994;61(5):935-40.
17. Ranganathan S, Ganguly AK, Datta K. Evidence for presence of hyaluronan binding protein on spermatozoa and its possible involvement in sperm function. *Mol Reprod Dev* 1994;38(1):69-76.
18. Ranganathan S, Bharadwaj A, Datta K. Hyaluronan mediates sperm motility by enhancing phosphorylation of proteins including hyaluronan binding protein. *Cell Mol Biol Res* 1995;41(5):467-76.
19. Nasr-Esfahani MH, Salehi M, Razavi S, Mardani M, Bahramian H, Steger K, et al. Effect of protamine-2 deficiency on ICSI outcome. *Reprod Biomed Online* 2004;9(6):652-8.