

Hard Ticks (Ixodidae) and Crimean-Congo Hemorrhagic Fever Virus in South West of Iran

Narges Sharifinia¹, Javad Rafinejad², Ahmad Ali Hanafi-Bojd², Sadegh Chinikar³, Norayer Piazak³,
Mojgan Baniardalani², Akbar Biglarian⁴, and Farhad Sharifinia⁵

¹ Department of Medical Entomology, School of Health, Ilam University of Medical Sciences, Ilam, Iran

² Department of Medical Entomology and Vector Control, School of Public Health, Tehran University of Medical Sciences, Tehran, Iran

³ Arboviruses and Viral Hemorrhagic Fevers Laboratory (National Reference Laboratory), Pasteur Institute of Iran, Tehran, Iran

⁴ Department of Biostatistics, University of Social Welfare and Rehabilitation Sciences (USWRS), Tehran, Iran

⁵ Veterinary Administration, DarrehShahr County, Iran

Received: 15 Oct. 2013; Accepted: 26 May. 2014

Abstract- Ticks are vectors of some important arthropod-borne diseases in both fields of veterinary and medicine, such as Lyme, tularemia, Rocky Mountain spotted fever, and some types of encephalitis as well as Crimean Congo hemorrhagic fever (CCHF). Iran is known as one of the main foci of CCHF in west of Asia. This study was conducted in DarrehShahr County because of the development of animal husbandry in this area to detect the fauna and viral infection of the hard ticks of livestock. A cross-sectional survey was conducted during 2011-2012 with random sampling in four villages. A sample of ticks was subjected to RT-PCR method for detection of viral infection. During the study period, 592 Ixodidae ticks were collected and identified as seven species of *Hyalomma asiaticum*, *Hy. marginatum*, *Hy. anatolicum*, *Hy. dromedarii*, *Hy. detritum*, *Rhipicephalus bursa* and *Rh. sanguineus*. More than 20% of these ticks were examined to detect the genome of CCHF virus while 6.6% were positive. All species of *Hyalomma* were found to be positive. A high rate of livestock was found to be infected with hard ticks, which can act as the vectors of the CCHF disease. Regarding infection of all five *Hyalomma* species captured in this area, this genus should be considered as the main vector of CCHF. Planning control program can be performed based on the obtained data on seasonal activity of Ixodidae to prevent animal infestation as well as to reduce the risk of CCHF transmission.

© 2015 Tehran University of Medical Sciences. All rights reserved.

Acta Medica Iranica, 2015;53(3):177-181.

Keywords: Hyalomma; Hemorrhagic fever, Crimean; Fever; Iran

Introduction

Ticks are blood-sucking arthropods; exist around the world as obligate parasites of the vertebrates (1). They transmit some viruses, bacteria, rickettsia and protozoa to human and animals so that it is a significant issue in the livestock industry (2). The environment is changing fast, human and livestock populations are increasing and therefore, patterns of relationship between ticks, hosts and parasites will change as a result of any variation in the density of each particle of this triangle (3). Knowledge about the epidemiology of vector-borne diseases, especially tick-borne diseases, will be useful in the control strategies for those diseases (4).

Crimean Congo Hemorrhagic Fever (CCHF) is one of the most common tick-borne diseases with a mortality rate of about 30%. This disease is endemic in large areas of Africa, as well as the eastern and central parts of the Europe. In recent years, the cases of the disease are significantly increased in Albania, Kosovo, Turkey and Iran while there are reports of CCHF from Greece (5). The epidemic form of CCHF usually occurs in areas where the public health services are poor, and high mortality may occur. Furthermore, climate, environmental and agricultural changes may affect the distribution of the ticks and the disease emergence (6). In the previous decade many cases of the disease have been reported from India, Russia, Eastern Europe,

Corresponding Author: A.A. Hanafi-Bojd

Department of Medical Entomology & Vector Control, School of Public Health, Tehran University of Medical Sciences, Tehran, Iran
Tel: +98 21 42933167, Fax: +98 21 88951393, E-mail address: aahanafibojd@tums.ac.ir

Hard Ticks and CCH

French, Pakistan, Afghanistan, Iran, Central Asia countries, Iraq, United Arab Emirates, Greece, Portugal, and different areas of the Africa continent (7).

Studies in Iran show the first report of CCHF virus in *Alveonatus lahorensis*, a soft tick collected in the north east of Iran (8). During 1998, two epidemics of the disease occurred in the eastern neighbors of Iran and led to death in some cases (7). The reports of the disease have submitted from some counties since 1999 (9). Although human infection and CCHF are recorded from 23 provinces of Iran, the most prevalence of the diseases is reported from Isfahan, Fars, Khorassan and Yazd provinces (10). Studies on the fauna of livestock ticks in relation to CCHF during 2000-2011 show 17 hard tick species are active in the studied areas as follows: *Boophilus annulatus*, *Dermacentor marginatus*, *Haemaphysalis punctata*, *Hae. sulcata*, *Hyalomma aegyptium*, *Hy. anatolicum anatolicum*, *Hy. anatolicum excavatum*, *Hy. asiaticum asiaticum*, *Hy. detritum*, *Hy. dromedarii*, *Hy. marginatum marginatum*, *Hy. schulzei*, *Hy. detritum*, *Ixodes ricinus*, *Rhipicephalus bursa*, *R. sanguineus* and *R. Turanicus* (11-24).

Because the Darreh Shahr county includes nomad people that, their main job is animal husbandry. Every year many animals are migrating to different parts of this area. Therefore, studying infection rate of livestock to ticks, tick fauna as well as the population at risk of CCHF seems to be necessary. Also survey on the infection rate of the collected ticks either introduce the vectors of CCHF in the area or is an important step toward improving planning for prevention of CCHF outbreak.

Materials and Methods

Ilam province is located in south west of Iran. This province is surrounded by Kermanshah province in the north, Lorestan province in the east, Khuzestan province in south and Iraq country in west. Darreh Shahr County with an area of 1480 km² and coordinates of 33°7'N and 47° 21' E, and elevation of 650 m above the sea level, is 142 km away from the Ilam city, the capital of the province (Fig. 1). This county has a tropical climate with long and hot summers and short and temperate winters. The highest and lowest temperatures in 2012 were recorded as 42 and -6 °C, respectively. Darreh Shahr County has a population of 56346. Nomads can be found in different areas of this county and animal husbandry is the main job of inhabitants.

This cross-sectional study was conducted during 2011-2012 in four seasons. Sampling was random in

four villages of the county: Gholamabad in north, Markazi in center, Farhadabad in south and Kolejooob in the east. These sampling sites were selected after consultation with the veterinary staff of the county.

Tick collection and virus detection

Sampling was carried out from all parts of the body of visited livestock (sheep, goat and cow). Collected ticks were put in the capped tubes, and all details were recorded, including weather information. The samples were then transferred to the laboratory for species identification (25). Regarding the type of the animal host, season and study village, a number of hard ticks were selected at random and RT-PCR were used to detect the viral infection (26). Sample size for this part of the study was calculated based on the infection rate of hard ticks in different parts of the country using the following equation.

$$n_0 = \frac{Z_{1-\alpha/2}^2 p(1-p)}{d^2},$$

where Z= 95%, P= 0.2 and d= 0.59.

SPSS 16, Spearman and χ^2 tests were used for statistical analysis of the data.

Results

During the period of tick sampling, 640 livestock were visited; 28.59% were found to be infected to hard ticks. A total of 592 hard ticks were collected from different parts of the infected animal bodies. Two genera of the hard ticks were identified: *Hyalomma* (90.21%) and *Rhipicephalus* (9.79%). The first genus included five species: *Hy. marginatum*, *Hy. anatolicum*, *Hy. asiaticum*, *Hy. dromedarii* and *Hy. detritum*; while two species were found to be related to the second genus: *Rh. sanguineus* and *Rh. bursa*. All species were collected from the four studied villages, although their frequency was variable in those villages in different seasons. Sex ratio of the collected specimens showed 82.1% females versus 17.9% males.

Seasonal activity of the hard tick was considered, and it was found that the highest activity is happening during spring and summer, whereas the lowest density of this arthropod observed in winter (Table 1). Statistical analysis showed a significant difference in the study seasons ($P<0.05$). None of the *Rhipicephalus* species was captured during autumn and spring visits.

Sheep was found to be the most infected animal host (59.6%), followed by goat (30.24%) and cow (10.13%). All species were found on all hosts, except for *Rh.*

bursa. There was no record of a collection of this species on cow (Table 2).

Out of 592 collected hard tick, 137 (23.1%) were selected at random for the study of CCHF virus infection. Among this sample 109 and 28 ticks were due

to *Hyalomma* and *Rhipicephalus* genera, respectively. Result of this survey confirmed viral infection in nine (6.6%) specimens. All infected species were due to the *Hyalomma* genus: *Hy. marginatum*, *Hy. dromedarii*, *Hy. asiaticum*, *Hy. anatolicum* and *Hy. detritum*.

Table 1. Frequency and prevalence of hard ticks of the study area in different seasons, Darreh Shahr County, West of Iran, 2011-2012

Tick species	Season								Total
	Spring		Summer		Autumn		Winter		
	No.	%	No.	%	No.	%	No.	%	
<i>Hy. anatolicum</i>	11	4.85	68	39.77	26	21.31	20	27.78	125
<i>Hy. asiaticum</i>	75	33.04	49	28.65	34	27.87	6	8.33	164
<i>Hy. detritum</i>	23	10.13	0	0	3	2.46	0	0	26
<i>Hy. dromedarii</i>	10	4.41	5	2.92	51	41.80	0	0	66
<i>Hy. marginatum</i>	57	25.11	42	24.56	8	6.56	46	63.89	153
<i>Rh. bursa</i>	14	6.17	1	0.58	0	0	0	0	15
<i>Rh. sanguineus</i>	37	16.30	6	3.51	0	0	0	0	43
Total	227	100	171	100	122	100	72	100	592

Table 2. Frequency and prevalence of hard ticks of the study area in different seasons, Darreh Shahr County, West of Iran, 2011-2012

Tick species	Animal host						Total
	Goat		Sheep		Cow		
	No.	%	No.	%	No.	%	
<i>Hy. anatolicum</i>	23	12.84	90	25.50	12	20	125
<i>Hy. asiaticum</i>	61	34.09	82	23.23	21	35	164
<i>Hy. detritum</i>	9	5.02	16	4.53	1	1.66	26
<i>Hy. dromedarii</i>	8	4.46	39	11.05	19	31.67	66
<i>Hy. marginatum</i>	43	24.05	105	29.75	5	8.34	153
<i>Rh. bursa</i>	11	6.14	4	1.13	0	0	15
<i>Rh. sanguineus</i>	24	13.40	17	4.82	2	3.33	43
Total	179	100	353	100	60	100	592

Discussion

Findings of this study show an infection rate of 28.59% in the livestock. This rate was higher than previous reports from several Iranian provinces, Ilam (19), Hamedan (12), Ghaemshahr (27), North West of Iran (28), and Azarbaijan-e-Sharghi (29); and lower from reports of Golestan (30), and some selected areas of Iran (31). The infection rates based on the animal host is consistent with the results of Sarani (2011) and Nasiri *et al.* (2009) that show the highest infection rate in sheep, goats and cattle, respectively (19,30).

Most of the collected ticks were species of the

Hyalomma genus. This is the same as previous studies in Ilam province (19,32), Yazd province (18) and Meshkinshahr (33). The dominant species varies in different studies. It indicates that the natural fauna of ticks is related to the ecological factors. The frequency of ticks is dependent on the season and climate, although in almost all previous studies, spring and summer had the highest infection rates, while winter is reported to have the lowest numbers (15,19,22,30,34-37). This may be the result of the change in the grazing pattern. So that in the spring and summer, when pastures have adequate forage and provide the food for livestock, host-seeking ticks have more chance to infect the livestock there.

Hard Ticks and CCH

Unlike, in the cold season, there is no favorable food in pastures and shepherds keep the flock in the fold, where they use acaricides for tick control.

Although minor differences exist between the results of this study and the surveys mentioned above, they can be justified due to difference in climate, weather, management and rearing of livestock farms and their plans for spraying and other tick control methods.

The highest infection rate of the tested ticks to CCHF virus was found to be in the spring and summer seasons. All infected ticks to CCHF virus were found to be species of *Hyalomma*. This genus had the highest infection rate in different parts of Iran, although some other species have also been infected (16,30,38-40). In Turkey the northwestern neighbor of Iran, it is found that *Rhipicephalus* ticks had more infection than *Hyalomma* (41). More than 90% of the collected ticks in this study were identified as *Hyalomma*, and so it is expected to find more infection rate in this genus. Of particular interest was finding the CCHF virus in all five collected species of *Hyalomma* in the study area.

Given that ticks are the most important vectors of CCHF and according to the results of this study, particular attention should be paid to combat these pests. In planning to control the ticks in the study area, it should be considered the effect of ticks on public health and economy of the community, sanitation, construction of anti-tick bath in the infected areas, organized distribution and using acaricides and training programs should be considered.

Acknowledgement

Thanks to everyone who contributed at different stages of this study, especially the staff of Arbovirus laboratory of Pasteur Institute of Iran, staff of veterinary organization of Darreh Shahr County, and Agriculture organization of Ilam province. This study is financially supported by Deputy of the Research, Tehran University of Medical Sciences.

References

1. Parola P, Raoult D. Ticks and tickborne bacterial diseases in humans: an emerging infectious threat. *Clin Infect Dis* 2001;32(6):897-928.
2. Vahedi Noori N, Rahbari S, Bokaei S. The seasonal activity of *Rhipicephalus bursa* in cattle in Amol (Northern Iran). *World Appl Sci J* 2012;18(4):590-3.
3. Hoogstral H, Valdez R, editors. Ticks (Ixodidae) from wild sheep and goats in Iran and medical and veterinary implications. 1st ed. Field Museum of Natural History 1980: p. 1-16.
4. Morzaria S, Katende J, Kairo A, et al. New methods for the diagnosis of *Babesia bigemina* the infection. *Mem Inst Oswaldo Cruz* 1992;87(Suppl 3):201-5.
5. Wolfel R, Paweska JT, Petersen N, et al. Low-density macroarray for rapid detection and identification of Crimean-Congo hemorrhagic fever virus. *J Clin Microbiol* 2009;47(4):1025-30.
6. Soares-weiser K, Thomass S, Gail G, et al. Ribavirin for Crimean-Congo hemorrhagic fever: systematic review and meta-analysis. *BMC Infect Dis* 2010;10:207.
7. Mardani M, Bijani B, Zare Kh. Case report: Histopathological findings in necropsy of a patient with fatal Crimean-Congo Hemorrhagic Fever in Iran. *J Res Med Sci* 2004;27(2):157-61.
8. Sureau P, Klein JM, Casals J, et al. Isolation of Thogoto, Wad Medani, Wanowrie and Crimean-Congo Hemorrhagic fever viruses from ticks of domestic animals in Iran. *Annals de Virologie* 1980;131(2):185-200.
9. Izadi Sh, Holakouie-Naeini K, Majdzadeh SR, et al. Prevalences of Congo-Crimean hemorrhagic fever in Sistan va Baluchestan Province: a serological study. *Payesh J* 2003;2(2):85-93.
10. Chinikar S, Ghiasi S, Moradi M, et al. Lessons of 10 years experience on CCHF in Iran. *BMC* 2011;5(Suppl 1):58.
11. Ranjbar Bahadori Sh. Study of species' diversity of animal ticks in Garmsar. *J Vet Res* 2003;58(1):11-4.
12. Moradi A, Telmadarraiy Z, Moradi A. Contamination rate of sheep to cattle ticks and its distribution in Bahar County. *J Vet Med* 2010;83(1):26-8.
13. Bahman Shabestari A, Davoodi J. Study on goat's tick fauna and seasonal variations of tick population in Zanjan Province. *Vet Res* 2009;5(2):79-84.
14. Rasooli S, Rajabi A, Jafari K, et al. Epidemiology of a severe pandemic could infect Sheep in Maragheh County. *Vet Res* 2011;10(4):61-6.
15. Hashemzadeh Farhang H, Khayatnoori M, Gharahdaghi Y, et al. Determined species diversity of hard ticks on referential sheep to the slaughter of Mahabad City. *Vet Res* 2011;12(4):61-6.
16. Telmadarraiy Z, Bahrami A, Vatandoost H. A Survey on fauna of ticks in West Azerbaijan Province, Iran. *Iran J Publ Health* 2004;33(4):65-9.
17. Asgarian F, Enayati A, Amooii A, et al. The fauna, geographic distribution and seasonal activity of hard Ticks from Sari Township in 2007-2008. *J Mazandaran Univ Med Sci* 2012;83(1):21-5.
18. Salim abadi Y, Telmadarraiy Z, Vatandoost H, et al. Hard ticks on domestic ruminants and their seasonal population dynamics in Yazd Province, Iran. *Iran J Arthropod Borne*

- Dis 2010;4(1):66-71.
19. Nasiri A, Telmadarraiy Z, Vatandoost H, et al. Tick infestation rate of sheep and their distribution in Abdanan County, Ilam Province, Iran, 2007-2008. *Iran J Arthropod Borne Dis* 2010;4(2):56-60.
 20. MotevalliHaghi F, Razmi G, Fakhar M, et al. The hard ticks (Ixodidae) fauna of livestock in Sari suburb, Northern Iran. *Comp Clin Pathol* 2013;22(1):5-8.
 21. Mirzaei Dehaghi M, Fathi S, NorouziAsl E, et al. Prevalence of Ixodid ticks on cattle and sheep southeast of Iran. *Trop Anim Health Prod* 2011;43(2):459-61.
 22. Nourollahi Fard SR, Fathi S, Norouzi Asl E, et al. Hard ticks on one-humped camel (*Camelus dromedarius*) and their seasonal population dynamics in southeast, Iran. *Trop Anim Health prod* 2012;44(1):197-200.
 23. Moshaverinia A, Dini M, Azizzadeh M. Prevalence of ixodid tick infestation of sheep in the Arasbaran region of Iran. *J Parasit Dis* 2012;36(2):230-33.
 24. Shemshad K, Rafinejad J, Kamali K, et al. Species diversity and geographic distribution of hard ticks (Acari: Ixodoidea: Ixodidae) infesting domestic ruminants, in Qazvin Province, Iran. *Parasitol Res* 2012;110(1):373-80.
 25. Walker AR, Bouattour A, editors. *Ticks of domestic animals in Africa, A guide to identification of species*. 1st ed. Winesconsin, U.K: Bioscience Reports; 2003: p. 157.
 26. Chinikar S, Persson SM, Johansson M, et al. Genetic analysis of Crimean-Congo hemorrhagic fever virus in Iran. *J Med Virol* 2004;73(3):404-11.
 27. Hosseini-vasoukolaei N, Telmadarraiy Z, Babamahmoodi F, et al. A survey of ixodid ticks parasiting domestic ruminants in Ghaemshahr District, Mazandaran Province, Iran. (Accessed in Jan 2014, 14, at <http://www.apjtm.net/admin/picture/UploadFile/20111231154156525.pdf>).
 28. Yakhchali M, Hosseini A. Prevalence and ectoparasites fauna of sheep and goats flocks in Urmia suburb, Iran. *Vet Arch* 2006;76(5):431-42.
 29. Telmadarraiy Z, Tighi S, Ghiasi SM, et al. Evaluation of Crimean-Congo hemorrhagic fever situation in East Azerbaijan Province: A serological and molecular Epidemiology Survey 2009. [EPUB ahead of print]
 30. Sarani M. Distribution of hard ticks and their infection to CCHF virus in Golestan Province using GIS method [Dissertation]. School Publ Health, Tehran Univ Med Sci., 2011.
 31. Rahbari S, Nabian S, Shayan P. Primary report on distribution of tick fauna in Iran. *J Parasitol Res* 2007;101(Suppl 2):S175-7.
 32. Gholami Parizad A. A survey on distribution of Ixodidae and Argasidae ticks in Ilam Province of Iran [Dissertation]. School Publ Health, Tehran Univ Med Sci., 1995.
 33. Tavakoli M. Survey on geographical distribution of ticks in Lorestan Province [Dissertation]. Tarbiat Modarres Univ., 1997.
 34. Khorramrooz A. Geographical distribution of Argasidae and Ixodidae ticks in Boyerahmad and Dena Counties, Kohgiluyeh va Boyerahmad Province of Iran. [Dissertation]. School Publ Health, Tehran Univ Med Sci., 2004.
 35. Rafieebarzoki M. Identification of hard ticks of domesticated ruminants in Semnan Province. *Pajouhesh Sazandegi* 2007;77(1):88-95.
 36. Salari Sh, Vatandoost H, Telmadarraiy Z, et al. Seasonal Activity of ticks and their importance in tick-borne infectious diseases in west Azerbaijan, Iran. *Iran J Arthropod Borne Dis* 2008;2(2):28-34.
 37. Tighi S. Survey on the Argasidae and Ixodidae ticks to CCHF virus by RT-PCR method and detection of its' IgG antibody in livestock and the at risk human cases in Bonad and Sarab Counties [Dissertation]. School Publ Health, Tehran Univ Med Sci., 2006.
 38. Telmadarraiy Z, Davari AR, Chinikar S, et al. Ruminant animal ticks and their role in CCHF transmission in Ghaen, South Khorassan Province, Iran during 2005. *Proceedings of the 11th International Congress of Parasitology*. Glasgow, Scotlandm, UK; 2006.
 39. Telmadarraiy Z, Moradi AR, Vatandoost H, et al. Crimean-Congo Hemorrhagic Fever: A seroepidemiological and molecular survey in Bahar, Hamadan Province of Iran. *Asian J Anim Vet Adv* 2008;3(5):321-7.
 40. Telmadarraiy Z, Ghiasi SM, Moradi M, et al. A survey of Crimean-Congo hemorrhagic fever in livestock and ticks in Ardabil province, Iran during 2004-2005. *Scand J Infect Dis* 2010;42(2):137-4.
 41. Tonbak S, Aktas M, Altay K, et al. Crimean-Congo hemorrhagic fever virus: genetic analysis and tick survey in Turkey. *J Clin Microbiol* 2006;44(11):4120-4.